



**UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460**

**OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES**

**MEMORANDUM**

**SUBJECT:** Analysis of Rodenticide Bait Use

**DATE:** September 10, 2004

**FROM:** Arnet Jones, Chief  
Biological Analysis Branch  
Biological and Economic Analysis Division

**TO:** Susan Lewis, Chief  
Reregistration Branch 1  
Special Review and Reregistration Division

Attached is the revised version of the Rodenticide Use Analysis document. This final version incorporates the comments received from three USDA offices (OPMP, APHIS, and FWS) and Stephen Frantz, who submitted comments in behalf of CDC. This version has cleared BEAD's Peer Review Panel.

## **Analysis of Rodenticide Bait Use**

### **Executive Summary**

This document presents an overview of the current use of nine rodenticide baits in the United States. Topics summarized include: the potential impact of rodents as disease vectors on human health; the damage caused by rodents to man-made structures and agriculture; a description of available market information, main use sites, target pests, and efficacy issues for these rodenticide baits; and alternative rodent control methods. The use profile for the nine rodenticides is summarized in Table 1 (P. 33).

The following major conclusions are put forward in this document:

- In the United States, rodents pose a major public health risk and cause economic damage to man-made structures and agriculture.
- Because rodenticide baits are an essential component of an integrated pest management approach to rodent control, this document concludes that the availability of rodenticides is necessary for the successful management of rodent populations.
- Since all nine rodenticide baits discussed in this document are currently registered for the control of commensal rodents in and around buildings, each of these is a potential alternative to each of the others in that use site situation.

Up-to-date data pertaining to amounts used, effectiveness, and prevalence of first generation anticoagulant resistance are not readily available. Section X (P. 33) lists topics for which new information will allow the Agency to refine this analysis.

### **I. Introduction**

For the past several years, EPA has been assessing the risks posed by rodenticides. As part of its ongoing work to reassess the safety of older pesticides, EPA issued a Reregistration Eligibility Decision (RED) for the Rodenticide Cluster in 1998. In that RED, EPA noted concern about potential adverse effects to birds and nontarget mammals, and announced a plan to further evaluate the potential ecological risks before issuing final decisions about reregistration eligibility. In 1999 EPA initiated a comparative ecological assessment, in which the Agency compared and ranked nine rodenticide active ingredients in terms of potential severity of risk. The comparative ecological assessment concludes that

there is adverse risk to nontarget organisms from all rodenticides, but certain compounds present more risk than others.

The statute under which EPA regulates pesticides, the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), requires consideration of benefits if adverse risks were assessed to be of concern. Thus, EPA must consider the benefits derived from the nine rodenticide products being evaluated before arriving at a decision regarding appropriate mitigation measures. The nine rodenticides included in the assessment are those addressed in the Rodenticide Cluster RED (brodifacoum, bromadiolone, bromethalin, chlorophacinone, diphacinone),<sup>1</sup> as well as zinc phosphide, warfarin, difethialone, and cholecalciferol.

The dual purpose for this document is, first, to summarize the available benefit/usage information in order to inform any risk management decisions, and, second, to request additional information not considered here (See Section X. Topics for Specific Public Comments.). EPA lacks complete usage information on rodenticide baits, particularly in the large homeowner market for rodenticides. Limited information is available concerning the professional applicator and agricultural markets. This document discusses the public health and other benefits associated with rodenticide products, briefly discusses efficacy, and concludes with a request for additional information on various aspects of rodenticide use and benefits. No attempt is made in this document to quantify the societal and environmental costs and benefits resulting from the use of rodenticide baits. This document, and its conclusions, will be revised, as appropriate, based on comments received during the 60-day public comment period.

## **II. Societal Benefits Derived from Rodent Control**

### **Background**

Many species of introduced and native rodents can cause health and economic damage to humans and their belongings. These species can transmit diseases; damage crops; consume and contaminate stored food supplies; disturb soil through burrowing activities; damage houses, other types of buildings and man-made structures; and prey on native species, including birds that nest on oceanic islands. It has been estimated that commensal rats cause between \$0.5 and \$1.0 billion of economic losses in the United States annually. This estimate was based on the assumption that there is one commensal rat per every two people in the country, at a time when the population of the United States numbered approximately 200 million, and that each rat consumes or damages between \$1 and \$10 worth of food and other materials, while contaminating 5 to 10 times more of it (Pratt et al., 1976). Commensal rats and mice have become adapted to live in close proximity to humans, thus having relatively easy access to almost unlimited food and shelter under certain conditions. Commensal

---

<sup>1</sup> In addition to brodifacoum, bromadiolone, bromethalin, chlorophacinone, diphacinone, the 1998 Rodenticide Cluster RED also presented EPA's reregistration decision for pival and its sodium salt, which were not supported and, therefore, determined to be ineligible for re-registration.

rodents are introduced Old World rats and mice in the family Muridae: the brown, sewer, or Norway rat (*Rattus norvegicus*), the black or roof rat (*R. rattus*), and the house mouse (*Mus musculus*). Some native rodents may also achieve a pest status comparable to commensal rodents and/or may be especially damaging to field and orchard crops or to turf. A high reproductive potential and mobility allows these rodents to rapidly fill available habitats and replace the individuals taken through available control methods.

Rodenticide baits, along with a variety of habitat modification and other managing techniques, are used to reduce the damage caused by native and introduced rodents, as well as other pest mammals. Rodenticide baits are especially useful for rapidly reducing rodent numbers in cases of major infestations. For commensal rodent control, rodenticide baits are best used within the context of an integrated pest management approach that emphasizes measures such as sanitation, exclusion, habitat modification, trapping, coordination at the community level, and public health education.

#### **A. Rodents and Human Health**

Rodenticide baits are used, sometimes as part of community rodent-control programs, in situations where major commensal rodent infestations must be reduced rapidly. Rodenticide baits often are used to control rodents which have entered homes and other buildings. Rodenticide baits are also used to eliminate rodents that remain after buildings have been rodentproofed (CDC 1999). The impact of coordinated rodenticide use can be significant in urban areas. For instance, during 1969-1981 the Center for Disease Control and Prevention (CDC)'s Urban Rat Control Program funded rat control projects in over 100 communities in 31 states. As a result, by 1980 there were 36,000 rat-free city blocks in the communities covered by the program, with another 20,000 blocks in the process of becoming rat-free. CDC estimates that by 1980 seven million people lived in areas that were made rat-free through this program (Blindauer, 1999).

Several New York City agencies are currently implementing coordinated rat control programs that include lot cleaning, cracking down on illegal garbage dumping, making regular inspections, and exterminating rats. The Department of Parks, for instance, uses an integrated pest management (IPM) approach, emphasizing prevention and exclusion methods that include frequent garbage collection, use of rodent-resistant closed containers, pruning of overgrown areas to reduce outside shelter, and masonry work to reduce rodent harborage inside structures. Rat populations are monitored to determine if the use of rodenticide baits is needed. This initiative was built on an IPM pilot project funded by CDC for implementation in a 48-block portion of Bushwick, Brooklyn, which had a long history of rodent infestations (Frieden, 2003).

Although in the past commensal rats and mice were the major cause of rodent-related public health concerns in the United States, the importance of white-footed and deer mice (*Peromyscus* spp.) as vectors of diseases, such as Lyme disease, human granulocytic ehrlichiosis, and hantavirus pulmonary syndrome, has increased in recent years.

## Disease transmission

**Salmonellosis** - Commensal rats and mice exposed to *Salmonella* bacteria in sewers or garbage may carry this pathogen in their gastrointestinal tract. Infected rats coming in contact with stored food, kitchenware, or food preparation surfaces may readily contaminate them with their droppings, which in turn could result in *Salmonella* food poisoning for humans exposed to the pathogens. Symptoms include nausea, diarrhea, and dehydration. This disease is rarely fatal (Blindauer, 1999).

**Plague** - In the Middle Ages, a plague pandemic transmitted to humans from infected rats by the bite of the oriental rat flea (*Xenopsylla cheopis*), was responsible for the death of about one third of Europe's population. Plague is caused by the bacterium *Yersinia pestis* and exists in three forms, systemic, pneumonic and bubonic. Symptoms include chills, generalized pain, and swollen lymph nodes. Untreated, the fatality rate could exceed 50% (Dept. Environ. Health, San Diego Co., undated). However, at present this disease is uncommon in humans in the United States, and no major urban plague outbreak has occurred in the country since 1924 (Schwarz, 2003). Plague can be treated with available antibiotics. During the 1980s, human plague cases in the United States averaged about 18 per year, with a fatality rate of one in seven (CDC, 2003). A plague reservoir exists in some wild rodent populations in several Western states, principally the rock squirrel, (*Spermophilus variegatus*) in the southwestern states and the California ground squirrel (*S. beecheyi*) in the Pacific states. Townzen, *et al* (1996) describe the management of a sylvatic plague epizootic in campgrounds in California. The primary rodent vector in that case was the California ground squirrel. Closing campgrounds, trapping fleas and squirrels to index their numbers and detect plague-positive individuals, insecticide dust for flea control, and Zinc Phosphide bait for controlling ground squirrels were elements of the program undertaken to combat the epizootic. Other rodents also identified as potential reservoirs include other ground squirrel species, prairie dogs (*Cynomys* spp.), wood rats (*Neotoma* spp.), chipmunks, and perhaps even deer mice and voles (CDC, 2003; Schwartz, 2003). Although cases of plague involving commensal rats have not been reported since 1925 (Dept. Environ. Health, San Diego Co., undated), there is a latent risk that as urban areas expand, the disease could be transmitted to commensal rats.

**Murine Typhus** - This disease, caused by *Rickettsia typhi*, is transmitted to humans by rat fleas when feces of infected fleas are scratched or rubbed into a flea bite area and into the bloodstream. Human cases of murine typhus increased rapidly in the United States from 1916 to 1945, with 21,572 cases being reported in 38 states from 1941 through 1945 (Andrews and Link, 1947). In 1944, 5276 cases were reported in eight southern States (AL, GA, FL, LA, MS, NC, TX, and TN). In Alabama in 1932-1933, the human death rate from murine typhus was 4.5 per 1000 reported cases, but nearly one-third of the cases involving people who were at least 65 years of age ended in a fatality (Andrews and Link, 1947). The incidence of murine typhus cases ultimately was greatly reduced through improved sanitation; insecticidal control of rat fleas; and lethal rodent control by use of rodenticide baits, chiefly

anticoagulants after they became available (e.g., Beall, 1946; Wiley, 1946; Hill and Morlan, 1948; Morlan and Hines, 1951; Mohr and Smith, 1957), and fumigants. The 10% DDT dust used to control rat fleas also produced some debility and mortality in rats (Dent, *et al* 1949).

Since disease management efforts took hold in the middle of the last century, murine typhus fever is relatively uncommon in the United States. Fewer than 50 cases are reported annually in the United States, mainly in California, Texas, and Hawaii (CDC, 2002). In some areas, such as southern California, the opossum (*Didelphis marsupialis*) can be a reservoir (Dept. Environ. Health, San Diego Co., undated). Several species of fleas have been implicated as vectors, including the rat flea (*X. cheopis*), the cat flea (on opossums) (*Ctenocephalides felis*), and the house mouse flea (*Leptopsylla segnis*). Although only a few cases are reported in Hawaii in typical years, in 2002 there were 47 cases recorded. Rodents suspected of being reservoirs in Hawaii include the Norway rat, the roof rat, the Polynesian rat (*Rattus exulans*) and the house mouse (CDC, 2002). Symptoms include fever, severe headache, general pain, and possibly a rash. It is rarely fatal at present (Blindauer, 1999).

**Rat Bite Fever** - Rat-bite fever is a bacterial disease caused by *Streptobacillus moniliformis* that can be acquired through the bite or scratch of a rodent or the ingestion of food or water contaminated with rat feces. Because cases are rarely reported in the United States, the true incidence of disease is unknown (CDC, 2003). Symptoms include influenza-like illness, rash, and arthritis, which may not be readily associated with a rat bite (Blindauer, 1999).

**Trichinosis** - Humans become infected with the nematode (an almost microscopic round worm) *Trichina spiralis* when ingesting improperly cooked infected pork. Nematode larvae parasitize the intestines and muscle tissue of humans. Pigs acquire the nematode by eating the carcasses of infected animals or grain food or garbage contaminated with feces from infected animals. Several wild and domestic mammals, including rats, cats, racoons, and bears, may serve as reservoirs for the pathogen, but the role of rats as a important reservoir may have been overemphasized in the past (USDA APHIS, 1972). According to the CDC, trichinosis infection in humans, once a common disease, is now relatively rare in United States. From 1991-1996, an annual average of 38 cases per year were reported (CDC, 2003). Symptoms include nausea, diarrhea, muscle aches, and may be fatal if the heart is involved (Blindauer, 1999).

**Leptospirosis** - This disease can be contracted by exposure to urine from infected commensal rats and mice. The causative organism, the spirochete bacterium *Leptospira icterohaemorrhagicae*, infect humans, dogs, and other domestic and wild animals by entering through mucous membranes or through cuts and scratches of the skin. Humans can become infected by handling contaminated items, from rodent bites, or from exposure to contaminated water. Symptoms due to infection may range from those associated with the common cold to kidney damage and liver failure (Blindauer, 1999).

**Rickettsial Pox** - This disease, caused by the obligate intracellular bacterium *Rickettsia akari*, can be transmitted to humans by the house mouse mite. Infection produces a chicken pox-like rash which is rarely fatal (Blindauer, 1999).

**Tropical Rat Mite** - In southern California and in the southern states, the tropical rat (*Ornithonyssus bacoti*) mite is a common ectoparasite of the roof rat. In rat-infested houses or buildings, this mite may also bite humans, causing irritation and dermatitis (Ebeling, 1975).

**Hantavirus** - A localized outbreak of hantavirus pulmonary syndrome resulting in several deaths occurred in 1993 in the area shared by Arizona, New Mexico, Colorado, and Utah, known as the Four Corners. The main virus host in this outbreak was the deer mouse (*Peromyscus maniculatus*). Since then, isolated cases have been reported in Louisiana, Florida, and New York involving the rice rat (*Oryzomys palustris*), the cotton rat (*Sigmodon hispidus*), and the white-footed mouse (*Peromyscus leucopus*) as reservoir hosts of several hantaviruses. The virus can be transmitted to humans through contact with accumulated urine and droppings from infected mice when tiny droplets of the material are stirred and become airborne (CDC, 1999). People may breathe in contaminated dust while cleaning or working in an infected area. Human exposure can also occur by introduction of the virus through the eyes, by ingesting contaminated food, or when bitten by an infected rodent. Symptoms include “fever, muscle ache, cough, rapid progression into severe lung disease” and often death (NPCA, 1994). Deaths occur due to rapid filling of lungs with bodily fluids (Larson and Morgenthaler, 1994). According to Boren and Valdez (2003), the mortality rate for humans is approximately 43% for the U.S. and 51% for New Mexico, while Mills *et al.* (2002) report that as of June, 2002, a total of 318 cases had been identified in 31 states, with a fatality rate of 37%. Hantaviruses do not cause illness in host rodents.

**Other Diseases** - Additional diseases known to be transmitted directly or indirectly by commensal and wild rodents in the United States include the following: lymphocytic choriomeningitis, toxoplasmosis, Colorado tick fever, Rocky Mountain spotted fever, Lyme disease, relapsing fever, babesiosis, western equine encephalitis, California encephalitis, human granulocytic ehrlichiosis, and cutaneous leishmaniasis, and *Yersinia pseudotuberculosis* (Blindauer, 1999).

## **B. Rodents and Domestic Animal Health**

Commensal rodents can also transmit diseases to domestic animals, such as leptospirosis and tapeworms to dogs and cats; brucellosis and foot-and-mouth disease to cattle; hog cholera and trichinosis to hogs; and salmonella, erysipelas, and fowl pox to poultry.

## **C. Rodents and Agriculture**

### **Preharvest**

Rodenticide baits are used extensively to manage rodents that feed on, contaminate, or cause various types of damage to a wide range of crops and farm infrastructure and equipment. Voles (*Microtus* spp.) can damage or kill young trees and shrubs by feeding on the inner bark layer of the trunks near the base, both above and below ground. When voles gnaw completely around the trunk (girdling), the flow of water and nutrients is disrupted and the tree or shrub is killed. In this way, voles damage fruit and nut orchards, windbreaks, and landscape plantings. Voles also feed on flower bulbs, on many types of garden plants and vegetables, and on various field and forage crops (Fisher and Hygnstrom, 1997; Salmon and Gorenzel, 1997; Montana Dept. Agr., undated). Deer mice can cause economic damage to Douglas fir and Ponderosa pine seeds in the West and Northwest. This is of concern in reforestation logged areas by direct seeding. Deer mice can also dig up planted seeds, including melon and alfalfa seeds. High populations of deer mice have caused damage to almond, avocado, citrus, pomegranate, and sugar beets.

Several species of pocket gophers (Geomyidae) feed on tubers and plants with succulent tap roots. Crops damaged include alfalfa, pastures, conifer plantations, row crops, and flower and vegetable gardens. Pocket gophers are a major problem wherever they occur in irrigated alfalfa. In Montana, for instance, yield losses in alfalfa may be up to 20-40%. In addition, pocket gopher mounds interfere with harvesting operations and damage harvest equipment. If gopher populations are moderate to dense, their mounds may cover 10-20% of the soil surface (Montana Dept. Agr., undated).

When locally abundant, the burrowing activities of ground squirrels and prairie dogs in agricultural areas can cause damage to harvesting equipment, pose a risk to livestock, and contribute to increases soil erosion. These rodents may also feed on grain crops.

Commensal rats and mice can be significant field pests of crops such as sugarcane, citrus, and Macadamia nuts. They can also damage agricultural equipment, including farm machinery, irrigation pipes and hoses.

### **Post-harvest**

Rodents often eat, damage, or contaminate (with urine, droppings, and hair) tons of stored food items, including grain, flour, cereals, sugar, vegetables, fruit, nuts, meat, animal feed, pet food, and any existing kind of stored edible material (Ebeling, 1975). The most damaging species in the U.S. are the commensal rodents and several species of voles. According to Hopf *et al.* (1976), cited by Brooks and Fiedler (1999), stored food losses to rodents in developing countries commonly range from 1% to 10%, but occasionally can be as high as 50%. Rodents typically contaminate far more stored grain

than they consume. During a year, a single mouse will deposit several thousand pellets (droppings) and about one pint of urine (Hygnstrom, 1995). The amount of rodent filth in stored grain is regulated by FDA and USDA: two or more rodent pellets or equivalent quantity of other animal filth per 1000 grams of grain reduces its quality to "U.S. Sample Grade," which can only be used for livestock feed (Brown, 1994; Hygnstrom, 1995). Although damage to stored food by rodents is not nearly as high in the U.S. as in developing countries, some losses do occur. The rodent damage to stored grain resulting from feeding and contamination was estimated at \$8.4 million per year in Nebraska alone (Hygnstrom, 1995). Such levels occur although efforts to mitigate rodent damage commonly are employed in the United States

#### **D. Structural/Industrial Rodent Damage**

Rodents' incisors never stop growing and are regularly worn down by their own self-wearing action and by gnawing hard materials (Frantz and Davis, 1991). In houses and buildings, commensal rodents can gnaw through gas pipes, electric wiring and its insulation, and building insulation, thereby creating a fire risk. They can also damage electronic and computer equipment. White-footed and deer mice often enter cabins and other buildings, where they may build nests and raise their young, causing damage to furniture, clothing, books, paper files, and other belongings (Timm and Howard, 1994). Damage from commensal rodents to farm buildings and equipment through gnawing, burrowing, and nest-building activities has been estimated at \$8.0 million per year (Hygnstrom, 1995)

#### **E. Predation on Island Species**

Introduced commensal rats have contributed to the endangerment and extinction of island plants and animals (Donlan *et al.* 2003). When commensal rats or other rodents become established on oceanic islands, they typically prey on the eggs, chicks, and sometimes the adults of ground- and hole-nesting birds, reptiles, and other types of organisms. The victims often include rare and endangered species. Often there are few or no native predators on such islands, and the resident birds have not evolved defense mechanisms which enable them to cope with introduced predators such as rats. If uncontrolled, this type of predation can lead to the local extinction of the affected bird populations (e.g., National Park Service, 2000).

#### **F. Other Types of Rodent Damage**

Rodents also cause economic losses to poultry operations by consuming feed, by causing structural damage to poultry facilities, and by vectoring pathogens, such as *Salmonella enteritidis*. New York poultry farmers, for instance, rank rodents, followed by flies, as the most serious pest problems affecting poultry farms (Harrington *et al.* 1998). Pocket gophers often damage lawns, golf courses, parks, and other noncrop areas through their burrowing activities. In Colorado alone, about three million acres of private lands are damaged to some extent by prairie dogs and other rodents

(State of Colorado Department of Agriculture, undated). Muskrats, nutria, and ground squirrels may damage levees, ditch banks, and culverts in agricultural areas.

### **III. Rodenticide Use**

There is little information publicly available on the use of and market for rodent control chemicals in residential, commercial, and agricultural settings. Data (U.S. EPA, 2004) that are available for residential and commercial use of rodenticide baits provide limited information on two broad markets for rodent control: the homeowner and the professional [or pest control operator (PCO)] markets. The private residential (“homeowner”) market includes products purchased by people to control rodents at home. The professional applicator market includes products purchased to control rodents in both residential and commercial settings.

The data available for agricultural rodenticide use are limited to general on-farm use and some specific field uses. A description of each market follows.

#### **A. Homeowner Market for Rodent Control**

Based on the available data, the annual value of the homeowner market for rodent control is estimated to equal more than \$90 million. The largest regional market is estimated to be the Southern United States (an estimated 35%-40% of the total market), followed by the Midwestern, Northeastern and Western United States. An estimated 90% of the market for the control of rodents is in the form of dry bait rodenticides, with other methods, such as glue boards, spring traps, and gas cartridges for burrowing rodents, making up the remaining 10% of the market.

D-Con® brand baits are the most widely used rodenticide product in the homeowner market. The most widely used active ingredient in d-Con® products is brodifacoum. Chlorophacinone, diphacinone, and other active ingredients also are available on the homeowner market.

#### **B. Professional (Pest Control Operator) Market for Rodent Control in Residential and Commercial Settings**

The estimated value of the market for rodent control products used by PCOs is more than \$15 million annually. The largest regional market for professional rodent control is the Southern United States (40% to 45% of the total market), followed by the Midwestern, Northeastern and Western United States. An estimated 75% to 80% of the professional market is rodenticide products, with the remaining 20% to 25% of the market including other methods of rodent control, such as glue boards and traps. An estimated 60% to 65% of professional rodent control is for house mice, 35% to 40% is for commensal rats, and 1% to 2% is for other rodents (e.g., pocket gophers, ground squirrels, etc.). An estimated 60% of the PCO work to control rodents is residential and 40% is commercial.

As much as 80% of the rodenticide sales to professional applicators is of products containing the active ingredients bromadiolone and brodifacoum. We estimate from available information that bromadiolone comprises 50% and brodifacoum 30% of the total professional market for rodenticides. The primary brand names for products containing bromadiolone are Contrac® and Maki®. For brodifacoum, the primary brand names are Talon®, Final® and WeatherBlok®. Other rodenticide baits with estimated use by professional applicators include difethialone, diphacinone, chlorophacinone, zinc phosphide, bromethalin and cholecalciferol.

### **C. Rodent Control in Agricultural Settings**

The available data (U.S. EPA, 2004; USDA National Agricultural Statistics Service, 1999) on the use of rodent control in agriculture estimate the use of rodenticide baits only. There is no information readily available on the extent of use of other methods, such as glue boards and traps, although it is assumed that these methods are used as well in agricultural settings. Based on the extent of use of these other methods in residential and commercial settings by homeowners and PCOs, it is assumed that the use of other methods in agricultural settings ranges from 10% to 25% of the total market for rodent control in these areas, with rodenticide baits accounting for the rest of the market.

According to available data, the primary rodenticide used in the field for rodent control is zinc phosphide. No other rodenticide appears to be used in significant amounts in agricultural fields. An estimate of the total amount of zinc phosphide used is not available, however, the data suggest that approximately 30% of total zinc phosphide use is in agricultural fields. Zinc phosphide is also used in noncrop areas (see below), in residential areas and on turf. An estimated 10% of its total use is in sugar beet fields, 10% of its use is in grain fields (e.g., wheat, barley, oats, etc.), and 10% of its use is on rangeland (U.S. EPA, 1998). The regional distribution of this use is not known.

The largest proportion of on-farm zinc phosphide use is in general farm areas. An estimated 40% of total zinc phosphide use is in and around farm buildings and structures, roads, ditches, and other noncrop areas (e.g., drainage ditches, irrigation canals, riding areas, feedlots, fence rows, etc.). In addition to zinc phosphide, the available data (USDA-NASS, 1999) provide an estimate for the use of brodifacoum, bromadiolone, bromethalin, cholecalciferol, difethialone, diphacinone, and warfarin in and around building structures; and for brodifacoum, bromadiolone and warfarin in roads, ditches and other non-cropland areas. Brodifacoum, bromethalin, diphacinone, and warfarin are also used in grain storage facilities for rodent control.

## **IV. Summary of Methods of Rodent Control**

### **A. Commensal Rodents in Urban and Suburban Settings**

In addition to the use of poison baits, there are a number of habitat modification approaches available for managing commensal rodent populations. These approaches aim at permanently denying

them of access to food, water, and shelter. Denying rodents access to food, water, and shelter is called “sanitation” (Bjornson and Wright, 1956). Denying rodents access to existing buildings is called “exclusion” (Frantz and Davis, 1991). Designing buildings to deny or deter rodent entry are called “proofing” and “stoppage,” respectively (Scott, 1982). Other types of habitat modification that may reduce the attractiveness of an area to rodents include changes to landscaping such as replacement of ground cover, such as English ivy with grass. To be successful in large, infested urban areas, these measures must be implemented on a community-wide basis. Most practical rodent control manuals, handbooks, and related literature include a section on nonchemical rodent management measures. A comprehensive review of this subject can be found in Frantz and Davis (1991).

Access to indoor and outdoor food sources can be limited by using rodentproof refuse containers, keeping garbage containers and bins tightly sealed, increasing the frequency of garbage pickup, rapidly and completely cleaning food spills, ensuring that no food or water remains exposed overnight at home, storing dried foods in rodentproof containers or rodentproof pantries, and retrieving seeds spilled from bird feeders on a daily basis. Eliminating non-toxic food sources also may make rodents more inclined to accept toxic baits.

Nesting and hiding places can be limited by thinning the planting density or by removing all the plant ground cover that offers protection to rodents near buildings, using plants that do not provide rodent harborage or attract rodents, removing or mowing tall vegetation growing in vacant lots; removing wood, construction debris, or any other materials stacked against buildings; cleaning up and reducing clutter in basements and other rarely-used rooms.

Young commensal rats can squeeze through openings “higher or wider than ½ inch” (Baker, *et al*, 1994). House mice can access openings ¼ inch by about 1½ inch (Frantz, 1988) or larger than ¼ inch (Baker, *et al*, 1994). To prevent rodents from gaining entry, buildings can be “rat-proofed” by sealing probable points of entry with heavy wire mesh or other appropriate materials. Potential points of entry may include air vents; openings around water and sewer pipes, electric lights, telephone wires, and TV cables; cracks around windows; doors and door frames, especially between lower edge and floor; holes in floors, walls, and ceilings; and any other type of opening that permits access to rodents. Trimming tree branches that overhang or touch homes or buildings keep roof rats, squirrels, and deer mice from using them to get access to homes.

Snap traps can be used to control rodents under low infestation levels, or where rodenticide use is too hazardous, or where the odor of dead rodents in inaccessible places would not be acceptable. Using snap traps correctly requires a good understanding of rodent habits. Glue boards are sticky material that trap rodents upon contact and work best with mice and immature rats. Naphthalene and paradichlorobenzene (PDB) may discourage rodents from entering an enclosed space that is not frequented by humans.

## **B. Native and Commensal Rodents in Agricultural/Field Settings**

Although poison baits are, in most cases, the fastest and most economical way of reducing pest rodent populations under agricultural or field conditions, several other rodent management methods exist, including the following:

- Keeping the orchard floor closely-mowed and maintaining weed-free strips under trees eliminates cover and reduces the chances of build-up of vole populations.
- Tree guards made of materials such as galvanized hardware cloth can be used to prevent tree girdling by meadow voles.
- On a small-scale basis, squirrels and chipmunks can be trapped using one of several kinds of live-traps available in the market.
- Ignitable gas cartridges and aluminum phosphide tablets are used to fumigate ground squirrel burrows. These materials are placed in burrows, which are then sealed with soil. Efficacy of fumigants is reduced in dry, porous soil.

## **V. Rodenticide Baits**

There are currently six anticoagulant rodenticides registered for use in the United States. These are: warfarin, chlorophacinone, diphacinone, brodifacoum, bromadiolone, and difethialone. The first three, often described as “first-generation anticoagulants,” generally require multiple feedings over several days to cause the death of a target rodent. Anticoagulants inhibit the formation of prothrombin, a key protein in the blood clotting process, thus leading to capillary damage and internal bleeding. Physiological resistance to first-generation anticoagulants has been detected in the United States and on other places around the world. According to Frantz and Madigan (1998), anticoagulant resistance may be counteracted with baiting schemes that alternate periods of baiting with warfarin, or other first generation anticoagulants, with periods of no anticoagulant exposure. This is a strategy that merits additional testing and validation.

Brodifacoum, bromadiolone, and difethialone are considered to be “second-generation anticoagulants.” The expression “second-generation anticoagulant” was coined to describe anticoagulant compounds that were believed to be effective against rodents that are resistant to the first-generation anticoagulants (Marsh, *et al*, 1980). Second-generation anticoagulants sometimes are also called “single-feeding anticoagulants” because rodents exposed to them may ingest a lethal quantity of the poison in a single night’s feeding. As the toxic symptoms of anticoagulant rodenticides do not take full behavioral effect for several days, however, rodents exposed to second-generation anticoagulant

baits may continue to feed relatively normally on them for 3 or 4 days and accumulate a “super-lethal dose.”

Anticoagulant rodenticide baits are used primarily to control commensal rats and mice and are mixed with grain products and other ingredients to make ready-to-use bait in meal, pelleted, or wax-block form. Meal and pelleted anticoagulant baits are intended for use in dry areas in and around buildings. Pelleted and meal formulations may be marketed in small plastic or paper bags (placepacks) which contain them at the site of placement until an animal breaks into the bag to feed on bait. Wax-block baits may be used where meal and pelleted baits are used and also may be used in sewers if the label of the specific product provides for such use. All placements of baits used to control commensal rodents in structural situations must be in tamper-resistant bait or stations if such placements otherwise would be accessible to children under six years-of-age, as well as to domestic animals and/or nontarget wildlife. Some commercial bait stations are designed especially to hold paraffinized bait blocks.

This assessment also considers rodenticides that are not anticoagulants. These compounds are zinc phosphide, cholecalciferol, and bromethalin. Zinc phosphide is an acute poison that may kill a target rodent as the result of a single bout of feeding. Zinc phosphide has a garlic-like smell that may attract some rodents and reportedly repel many nontarget species. Zinc phosphide’s natural emetic action may, in some cases, serve to protect some nontarget species from toxicosis after they have eaten bait containing this active ingredient (Cornell University, 2001). Exposure of target rodents to untreated grain (prebaiting) is generally recommended to increase bait acceptance and is required for some use patterns. Bromethalin induces anorexia after a toxic, and usually fatal, amount has been consumed. Death ensues within one to several days after consumption. Cholecalciferol may require several feedings for death to occur.

Although all nine rodenticides briefly discussed above are registered for commensal rodent control, only zinc phosphide, chlorophacinone, diphacinone, and warfarin are registered for control of small native mammals, including “field” rodents such as voles, deer mice, ground squirrels, pocket gophers, prairie dogs, and kangaroo rats; jack rabbits; and moles. Target species claims vary according to chemical and label. Depending on application directions and use restrictions, baits used to control native rodents in agricultural and nonagricultural lands can be broadcasted or applied by hand in selected areas.

#### **A. Warfarin and its Sodium Salt**

Warfarin, the first anticoagulant rodenticide to be developed, was first registered in 1950. Warfarin was named after the foundation that developed it, the Wisconsin Alumni Research Foundation (WARF) (Link, 1959). EPA issued a Registration Standard for Warfarin in 1981 (EPA, 1981) and issued the Warfarin RED in 1991. Activities associated with those reregistration efforts brought the Warfarin products that remained registered up to the standards of the times and led to the cancellation of many other Warfarin products. Although resistance to warfarin and other first

generation anticoagulants has been detected in rats and mice in Europe since the late 1950s, current reliable estimates of its prevalence in the United States are not available.

Warfarin is formulated predominantly as dry (meal, pelleted, or paraffinized) baits. All but one of these baits are 0.025% active ingredient, with the exception being a 0.054% warfarin bait registered only for control of house mice. Other registered end-use products containing warfarin include 0.3% and 0.5% active ingredient concentrates for preparing baits. The sole remaining registered formulation of sodium salt of warfarin (0.54% active ingredient) is designed to be diluted with water to make liquid baits.

Warfarin is federally registered mainly for control of Norway rats, roof rats, and house mice in and around homes, agricultural buildings, and commercial and industrial sites, including food and feed handling establishments. Warfarin is also registered for deer mouse and white-footed mouse control in and around homes, agricultural and industrial buildings and similar structure; in parks, woodlots, yards and lots surrounding residential buildings and noncrop areas near agricultural buildings. Finally, there is a recent warfarin registration for the control of several species of moles (eastern, star-nose, hairy-tailed, coast, broad-footed, Townsend) on lawns, turf areas, golf courses, and other non-food grassy areas. The warfarin bait registered for mole control must be applied directly into their underground tunnels.

## **B. Chlorophacinone**

Chlorophacinone is a first-generation anticoagulant that was first registered in 1971. Chlorophacinone may kill some rodents after a single night's feeding, but multiple feedings are needed in most cases. Death in rodents occurs within three to ten days.

Chlorophacinone is formulated as tracking powder, loose-grain bait, paraffinized pellets, bait in ready-to-use placepacks, and paraffin blocks. Bait formulations contain 0.005% active ingredient. Baits are applied as often as needed for controlling commensal rodents. Most field uses have a limited number of applications. Both restricted use and unclassified<sup>2</sup> products are registered.

Most of the unclassified products are ready-to-use baits registered for structural use to control commensal rodents. Application rates for placepacks and paraffinized blocks are similar to those for loose baits, but application amounts typically are expressed in terms of the numbers of placepacks, blocks, or pieces of blocks to deploy at each placement location. There also is one 0.005% chlorophacinone bait registered to control pocket gophers and another registered to control moles. These unclassified products must be applied directly into the underground burrow systems of pocket gophers and moles, respectively.

---

<sup>2</sup> The U.S. EPA does not normally classify products for general use; products that are not restricted remain unclassified. See 40 CFR 152.160(a).

The restricted-use products containing chlorophacinone include 0.2% a.i. tracking powders that are federally registered for indoor use to control commensal rodents and some 0.005% and 0.01% a.i. baits registered under Section 24(c) of FIFRA to control various types of field rodents. At the time of product reregistration, all chlorophacinone products registered for above-ground uses to control field rodents will be classified as Restricted Use Pesticides.

Chlorophacinone is federally registered for control of commensal rodents in and around homes; industrial, commercial, or agricultural buildings and structures; and inside sewers. There also are 24(c) “special local needs” (SLN) registrations authorizing use of chlorophacinone baits in 19 states for control of numerous small mammals in a variety of agricultural and non-agricultural field sites. Chlorophacinone is registered for control of eastern moles (*Scalopus aquaticus*), star-nosed moles (*Condylura cristata*), or moles (*Scapanus* spp.) in lawns, golf courses, and other turf areas; and pocket gophers in lawns, golf courses, range land, and noncrop areas. It is also registered to control various types of voles (*Microtus* spp.) in dormant and non-bearing orchards; California voles in artichoke fields; and ground squirrels around farm buildings, in orchards, alfalfa, rangelands, pastures, and noncrop areas. Other target mammals claimed on SLN labels for chlorophacinone baits are deer mice (*Peromyscus maniculatus*), wood rats (*Neotoma*), chipmunks (*Eutamias* spp.), prairie dogs (*Cynomys* spp.), muskrats (*Ondatra zibethica*), nutria (*Myocastor coypus*), and jackrabbits (*Lepus* spp.). Field use sites listed on labels include orchards, forests, groves, nurseries, tree plantations, inside transport vehicles, commercial transportation facilities, and food processing, handling and storage areas and facilities.

### **C. Diphacinone and its Sodium Salt**

**Background Information** - Diphacinone is a first-generation anticoagulant rodenticide first registered in 1960. As with chlorophacinone, diphacinone may kill some rodents after a single night’s feeding, but multiple feedings will occur and may be needed to cause death.

Diphacinone is federally registered for control of commensal rodents in and around homes, industrial and agricultural buildings and similar man-made structures; in sewers; in wet or damp areas including river banks, irrigation ditches, gullies, railroad tracks, loading areas, along fences, under granaries, garbage dumps, and landfills. Diphacinone is also federally registered for control of pocket gophers (*Thomomys bottae*, *T. talpoides*, *T. mazama*, *T. townsendi*, *T. bulbivorus*, *T. monticola*, and *Geomys bursarius*) in rangeland, grain fields, forage crops, hay and alfalfa crops, vegetable crops, forest, nurseries, and noncrop areas, including parks and around homes.

Twenty-two states currently have at least one 24(c) special local needs diphacinone registration for field uses. Uses of these products include control of voles, mainly *Microtus* spp., in dormant bearing and/or non-bearing tree fruit orchards in New Hampshire, Ohio, Pennsylvania, Virginia, Washington, West Virginia; control of voles in orchards, Christmas tree farms, commercial nurseries, and tree plantations in North Carolina; control of ground squirrels (*Spermophilus* spp.) and muskrats in

noncrop areas, control of jack rabbits (*Lepus californicus*) in noncrop areas bordering agricultural fields and at airports, control of California voles (*Microtus californicus*) and montane voles (*M. montanus*) in orchards and groves, and control of wood rats (*Neotoma* spp.) in/around cabins and plantations of citrus trees and conifers in California; control of ground squirrels (*Spermophilus* spp.) in levee or ditch banks, around farm buildings, along fence lines, in orchards, in or near crops, and in noncrop areas in Washington (State); control of voles around small grain crops in Washington and Idaho; control of cotton rats (*Sigmodon hispidus*), rice rats (*Oryzomys palustris*) and Florida water rats (*Neofiber alleni*) in noncrop areas adjacent to sugar cane and sweet corn fields in Florida; control of mongoose in Hawaii and the Virgin Islands for protection of numerous species of ground nesting birds, including endangered bird species; control of rats and mice around the perimeter of planted fields and nurseries in Florida; control of commensal rats in forests, offshore islands and other noncrop outdoor areas in Hawaii and the Virgin Islands; control of commensal rats for conservation purposes on islands in the Alaska Maritime National Wildlife Refuge; and control of deer mice, jackrabbits, chipmunks, muskrats, woodrats, voles.

The only remaining registered products containing sodium salt of diphacinone as the active ingredient (at 0.106%) are registered only for mixing liquid baits to be used to control commensal rodents in and around homes, industrial and agricultural buildings, and similar man-made structures.

#### **D. Brodifacoum**

Brodifacoum, first registered in 1979, is a second-generation anticoagulant and the most widely used by homeowners in urban areas at present. A single night's feeding may be sufficient to kill a target rodent within a three to ten days, but rodents often feed and behave normally for 2-3 days after their first exposure to the bait.

Brodifacoum is formulated as meal baits, paraffinized pellets, and paraffin bait blocks. All end use formulations contain 0.005% active ingredient. Baits may be applied as often as necessary. All registered brodifacoum products are unclassified.

Brodifacoum is currently registered for use only against commensal rats and mice in and around homes, agricultural buildings, commercial buildings, public buildings, and industrial buildings, inside transport vehicles, in and around related port or terminal buildings, in alleys, and in sewers.

At times, Brodifacoum has been used under §18 emergency exemptions from the requirements of FIFRA in Federal conservation programs to control rodents on oceanic islands. However, Brodifacoum does not have a Section 3 registration for that purpose.

### **E. Bromadiolone**

Bromadiolone is a second-generation anticoagulant which first was registered in 1980. Bromadiolone is formulated as meal baits, paraffinized pellets, and paraffin bait blocks. All formulations contain 0.005% a.i. Some pelleted baits and meal baits are sold in placepacks.

Bromadiolone is registered for control of commensal rats and mice in and around buildings, inside transport vehicles, in alleys, and in sewers in urban areas. It is limited to indoor use in homes and agricultural buildings in non-urban areas.

### **F. Difethialone**

Difethialone is a second generation anticoagulant rodenticide that was first registered in 1995. Difethialone is registered for the control of Norway rats, roof rats, and house mice in and around homes and industrial, commercial, and public buildings in urban areas; in transport and cargo vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. In non-urban areas, difethialone may only be used inside of homes and agricultural buildings.

### **G. Zinc Phosphide**

Zinc phosphide was registered in the United States in 1947, the first year of the Federal pesticide registration program under FIFRA. Prior to that time, zinc phosphide was used as a rodenticide in the United States and was used in that capacity in Italy as early as 1911. Once ingested, zinc phosphide reacts with moisture in the gastrointestinal tract to liberate phosphine gas, which is the lethal agent.

Zinc phosphide is the rodenticide most commonly used in agriculture. It is available in ready-to-use dry baits (whole-grains and pellets) and in the form of a dry concentrate to be used by applicators to prepare toxic baits. A 10% zinc phosphide tracking powder is registered to control house mice. All zinc phosphide end-use products are classified as Restricted Use Pesticides except those that are 2% active-ingredient or less and are limited to use in and around buildings (e.g., to control commensal rodents) or to manual, subterranean applications to the burrow systems of pocket gophers or moles.

Depending upon the specific use pattern and label, zinc phosphide baits may be applied manually, by ground or aerial bait broadcasting equipment, or by use of other specialized equipment such as trail builders or burrow builders. As zinc phosphide baits often are not readily accepted by target rodents, labels often advise that bait-acceptance tests be conducted prior to baiting operations. Depending upon the specific use pattern, labels may recommend or require prebaiting the entire infested area with rodenticide-free grain of the type to be used in the toxic bait to accustom target rodents to feeding on that particular grain.

Zinc phosphide is registered for control of commensal rats and mice, white-footed mice (*Peromyscus leucopus*), and voles in and around homes and industrial, commercial, agricultural, and public buildings. It is also federally registered for use against a wide range of small mammals, mainly rodents, such as the following: meadow voles in alfalfa and timothy hayfields; muskrats and nutria in areas where they damage levees, irrigation ditch banks, or water impoundments, or where they might feed on rice, soybean, milo, corn, or damage turf; pocket gophers in rangeland and Christmas tree plantations; pocket gophers and moles in grain fields, forage crops, hay and alfalfa crops, and vegetable crops, forest areas, parks, nurseries, lawns, golf courses, homes, and other noncrop areas; voles, white-footed and deer mice, and jumping mice (*Zapus* spp.) in grape vineyards; voles and white-footed mice in berry production areas; voles in sugar beets; voles and deer mice in noncrop areas, including right-of-ways, lawns, parks, nurseries, and golf courses; voles and white-footed mice in pastures, ornamentals, orchards, vineyards, rangelands, forests, reforestation areas, lawns, golf courses, parks, nurseries, and highway medians; roof rats in macadamia nut orchards; deer mice and voles in orchards, groves, nursery stock (ornamental and non-bearing fruit trees), and conifer and hardwood seeding plantations; prairie dogs and ground squirrels in rangeland and reforestation seedlings and plantings; kangaroo rats (*Dipodomys* spp.) in rangeland, pastures, and noncrop areas; rats (Polynesian, Norway, roof, rice, Florida water, cotton) in sugarcane fields; California ground squirrel and voles in noncrop rights-of-way; voles; ground squirrels in dormant orchards or vineyards, ornamentals, cemeteries, golf courses, nurseries, canals and ditch banks, rangelands, pastures, lawn and turf grasses, and along fence rows; prairie dogs on rangelands, pastures, and reforestation seedlings and plantings in the western United States; woodrats in rangelands, pastures, noncrop right-of-ways, dormant orchards, tree farms, and rural agricultural buildings.

#### **H. Cholecalciferol**

Cholecalciferol, also known as Vitamin D<sub>3</sub>, was first registered in 1984. Cholecalciferol triggers the mobilization of calcium from the bone matrix into blood plasma, resulting in death from hypercalcemia in 3-4 days after ingestion of a lethal dose. Cholecalciferol is not an anticoagulant. Cholecalciferol is formulated into baits containing 0.075% active ingredient.

Cholecalciferol is registered for the control of Norway rats, roof rats, and house mice in and around homes; industrial and commercial buildings; and similar man-made structures, inside transport and cargo vehicles (ships, trains, aircraft), and in related port or terminal buildings.

#### **I. Bromethalin**

Bromethalin is a diphenylamine rodenticide that acts, after one or more feedings, by blocking nerve impulse transmission, causing paralysis of the central nervous system and respiratory arrest in 2-4 days. Bromethalin was initially registered in 1982.

Bromethalin is formulated into various types of rodenticide baits, including paraffinized blocks, meal baits, and pelleted baits. Most formulated products contain 0.01% active ingredient. Some bromethalin baits are sold in placepacks. Two products consist of nine 0.01% bromethalin bait in a ready-to-use bait station. These bait stations have not been shown to be tamper-resistant.

Bromethalin is registered for control of commensal rats and mice in and around homes, industrial and agricultural buildings, and similar man-made structures; in alleys, in transport vehicles (ships, trains, and aircraft), and in and around related port or terminal buildings.

## **VI. Efficacy Considerations**

The rodenticide active ingredients discussed in this document were subjected to the product performance (efficacy) data requirements that were in effect at the time that they were registered. Those requirements have changed over time, with the most significant developments being the amendments to the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) passed 1972 and 1978, the development of the Product Performance portion (Subdivision G) of the *Pesticide Assessment Guidelines* (Schneider and Hitch, 1982), and the promulgation of the regulations published in 40 CFR §158.640.

The 1972 amendments to FIFRA required EPA to formalize its data and testing requirements through publication of relevant guidelines, instituted data compensation requirements, and required EPA to “register and reclassify pesticides registered” under FIFRA prior to the effective date of “regulations promulgated” to provide for “registration and reclassification of pesticides under the provisions of” FIFRA, as amended. This “registration and reclassification” activity later came to be known as “reregistration”.

The 1978 amendments authorized conditional registration of pesticide products, expanded on the subject of reregistration, and granted discretionary authority to the EPA administrator to “waive data requirements pertaining to efficacy.” EPA thus initiated a policy of waiving efficacy data requirements for all pesticidal claims except those pertaining to control of pests considered to be of significance to public health. The efficacy waiver policy was expanded in 1982 to include essentially all pesticidal claims. The policy soon was changed back to one of routinely requiring efficacy data to support claims for controlling pests of significance to public health and emphasizing that the waiver policy only applied to the efficacy data requirement. Registrants were expected to test pesticide products to verify that they were efficacious. The efficacy data submission requirements and policies are summarized in 40 CFR §158.640.

Of the claims being made for the rodenticides discussed in this document, the following rodent types have been considered to be of significance to public health: Norway rats, roof rats, house mice, *Peromyscus* spp. mice (except for endangered subspecies), ground squirrels (*Spermophilus* spp.), and prairie dogs (*Cynomys* spp., except for the endangered Utah prairie dog, *C. parvidens*).

Applicable efficacy data must be submitted or cited to support claims for controlling these species, which are considered to be of public health significance. Distinctions are not drawn between urban and rural, domestic and industrial, or structural and agricultural use sites because rodent-vectoring diseases may be spread to humans in any of these locations. There is a public health aspect to virtually all efforts to control these species.

As indicated in 40 CFR §158.640, EPA reserves the right to require that efficacy data be submitted “on a case-by-case basis for any pesticide product.” For the rodenticides discussed here, that authority has been considered in a few cases in which the claims being evaluated seemed unlikely for the product formulation involved.

Four of the nine rodenticides discussed in this document were registered prior to passage of the 1972 amendments. In order of the date of initial registration, those compounds are: zinc phosphide, warfarin, diphacinone, and chlorphacinone. In addition to the efficacy data that may have been required for registration, baits containing these active ingredients -- especially zinc phosphide and warfarin -- were tested rather extensively for effectiveness against commensal rodents under actual use conditions by university, municipal, and public health personnel (e.g., Emlen and Stokes, 1947; Hayes and Gaines, 1950). Additional uses of these four compounds that were accepted under Section 3 of FIFRA after passage of the 1972 amendments were subject to prevailing data requirements and, if efficacy data submission requirements applied, were thoroughly researched (Tietjen, 1976; Tietjen, 1979; Tietjen and Matschke, 1981).

The other 5 rodenticides considered here (brodifacoum, bromadiolone, bromethalin, cholecalciferol, and difethialone) were not registered until after passage of the 1978 amendments. These 5 compounds were extensively tested for efficacy according to procedures similar to those described in Subdivision G of the *Pesticide Assessment Guidelines* with respect to claims for controlling commensal rodents. Such testing entailed laboratory assessments of toxicity to target species, laboratory “choice” feeding trials to assess palatability and lethality of candidate bait formulations, and regional field testing. Although Bromethalin was initially registered in 1982, during the time of the expanded efficacy waiver, its laboratory and field testing regimens were largely completed before that policy went into effect.

Research has shown that seemingly small changes to bait composition can have drastic adverse effects upon palatability to commensal rodents. Therefore, specific bait formulations claimed to control commensal rodents are required to be screened for efficacy. Most commensal rodenticide bait formulations registered since the 1972 amendments have been screened for effectiveness against Norway rats and house mice, if both species are claimed on product labels. A few such products registered during the period of the expanded efficacy waiver (mid 1982 to early 1984) were not screened for efficacy prior to registration, but will be if they are to be reregistered. Warfarin baits registered prior to the 1972 amendment were screened for efficacy prior to their being reregistered in

the early 1990's. Registered commensal rodenticide baits must be screened for efficacy again if they are reformulated.

Cholecalciferol and difethialone were registered after November 1, 1984 and, therefore, are not subject to reregistration requirements. Prior to registration, however, these active ingredients were thoroughly tested for efficacy against commensal rodents in laboratory and field trials. All registered commensal rodenticide bait formulations containing these compounds have been screened for efficacy.

For registered baits products containing the compounds currently undergoing registration (zinc phosphide, diphacinone, chlorophacinone, brodifacoum, bromadiolone, and difethialone), new efficacy data generally will be needed for reregistration only if their registrants: (1) seek to modify existing bait formulations; (2) desire to make claims of effectiveness which go beyond simple claims for controlling the target species claimed on the products' labels, or (3) have not yet supported some claims for which submission of efficacy data is required.

Although the methods for screening rodenticide baits for efficacy in laboratory tests vary somewhat according to the type of product and the nature of the claim being investigated, the product performance criteria are similar across the nine compounds being considered in this document. In controlled laboratory tests with fresh "dry" or liquid baits claimed to control commensal rodents, at least 90% of the rodents exposed to the rodenticide must die for the bait to be considered to be efficacious. For multiple-feeding anticoagulants (e.g., warfarin, diphacinone, and chlorophacinone), there also is a requirement that fresh baits be accepted as 33% or more of total intake in a choice test involving EPA's standard challenge diet for dry baits or water for liquid baits. Slightly lower criteria (80% mortality and 25% acceptance) apply when experimentally "weathered" anticoagulant baits are evaluated for suitability for use in wet or damp areas (e.g., sewers) or are claimed to be "weather-resistant."

The performance criteria in field tests are similar across active ingredients used in rodenticide baits. EPA accepts field efficacy studies of rodenticide baits if appropriate procedures were employed in the research and if rodent activity indices indicate that at least 70% control of the baited population was obtained. All the rodenticide compounds discussed here have been tested according to essentially similar criteria and are considered by EPA to be useful for controlling commensal rodents.

Three of the nine compounds under discussion are registered for use to control "field" rodents of public health significance (e.g., ground squirrels, prairie dogs, *Peromyscus* spp.). These compounds are: zinc phosphide, diphacinone, and chlorophacinone. Zinc phosphide bait products labeled for controlling these species and other field rodents (e.g., *Microtus* spp. voles; *Geomys* and *Thomomys* spp. pocket gophers; non-endangered kangaroo rats, *Dipodomys* spp. etc.) typically are registered under Section 3 of FIFRA. Most diphacinone and chlorophacinone products labeled for field rodent uses are available only in certain States under "special local needs" registrations issued according to the provisions of Section 24(c) of FIFRA. Reregistration of field uses for these three compounds will entail

submitting or citing relevant data, including efficacy data regarding claims for controlling pests that are of significance to public-health claims. Some information of that nature that has been reviewed to date is tacitly considered in this assessment. New efficacy studies may be needed if there are any public-health claims that are not fully supported by existing data.

Although rodenticide compounds registered for similar uses have been required to meet essentially the same performance criteria, the field tests that have been reported to EPA to fulfill efficacy data requirements for these chemicals were run at different sites, at different times, and often by different personnel. That these and other factors may affect study outcomes should be kept in mind when comparing results from such tests.

Relatively few field efficacy trials have been conducted specifically to compare the performance of various rodenticide baits under United States field conditions. Generally, such trials were conducted under conditions in which control of the target species had proven to be difficult. For example, Tickes, *et al* (1982) ran comparative efficacy trials with valley pocket gophers using three compounds that are registered for that purpose, but reported that none of the six bait formulations tested appeared to control that organism well. Some of the comparative studies that have been reported have primarily involved use of compounds which are not included in this evaluation and/or are no longer are registered (e.g., Albert and Record, 1979; Emlen and Stokes, 1947).

Ashton, *et al* (1983) conducted comparative efficacy trials for several currently registered rodenticide baits containing warfarin, bromethalin, brodifacoum, or bromadiolone. Baits were evaluated for efficacy against house mice and Norway rats. As those trials were performed at sites where resistance to warfarin (see below) had been detected, the test results obtained for warfarin were likely lower than typical results would have been with populations of non-warfarin resistant rodents.

## **VII. Factors Affecting Rodenticide Bait Selection**

Although the nine rodenticide compounds considered here have been required to meet similar criteria in efficacy trials, several additional factors must be considered when assessing the degree to which compounds registered for similar uses may serve as alternatives to one another. These factors are identified and briefly discussed below.

**Registered Uses** - Rodenticide baits may only be used at the use sites that are indicated on their labels and may not be applied by application methods that are prohibited by label text (e.g., “Do not broadcast bait”, “Apply this product only by the methods prescribed on this label”, “This product must be applied directly into pocket gophers’ underground burrow systems”, etc.). While all of the nine rodenticides considered here are registered for controlling commensal rodents, zinc phosphide is the only one that has been accepted under Section 3 of FIFRA for broad usage to control field rodents and is the only one for which tolerances have been issued to allow for food and feed uses. These circumstances leave users with no toxic-bait alternative to zinc phosphide for many uses. In certain

states, baits containing Chlorophacinone or Diphacinone are available under “special local needs” [Section 24(c)] registrations for control of field rodents.

**Classification** - Not all rodenticide baits are available to all prospective users. Products labeled as “Restricted Use Pesticides” may only be purchased by certified applicators and may only be used by such applicators or by persons under their direct supervision. Such products include zinc phosphide baits registered for above-ground uses to control field rodents. All above-ground uses of any of the other eight compounds to control field rodents will likely be classified as restricted use pesticides at the time of reregistration.

**Speed of Kill** - How fast a compound kills a given target species may affect the choice of the toxicant. Of the nine rodenticides considered, zinc phosphide and bromethalin come closest to being true “acute” rodenticides, with individual rodents generally feeding on the bait for only one day and dying within one or two days after feeding occurs. With cholecalciferol baits, significant feeding usually occurs on more than one day with death ensuing several days later.

Anticoagulants generally take at least four days from the onset of feeding until rodents begin to die. It may take two weeks or more for some individuals to die. Although second-generation anticoagulants also have been touted as “single-feeding” anticoagulants, that expression is misleading. It has been shown that although the amount of bait containing bromadiolone, brodifacoum, or difethialone eaten in one 24-hour period can be sufficient to cause the deaths of rodents, the animals generally continue to feed normally for two or three more days. Thus, free-ranging rodents exposed to a palatable second-generation anticoagulant bait would be expected to eat nearly as much of it as they would if exposed to a palatable first-generation anticoagulant bait, thereby accumulating a “super-lethal dose.”

**Perceived risks to nontarget species** - The extent of the applicator’s knowledge and concern about nontarget effects would likely play a part in the selection of a toxicant. If, for example, bait applications are to be primarily used in indoor locations to which children or nontarget animals, such as pets and livestock, do not have access, an applicator might feel more comfortable in using a second-generation anticoagulant than if outdoor placements were needed in areas frequented by birds or nontarget mammals.

**Resistance** - Resistance to first-generation anticoagulants has been detected in commensal rodent populations in many places throughout the world including some localities in the United States (Frantz and Padula, 1980; Frantz and Madigan, 1998). Efficacy data submitted to the Agency for brodifacoum indicate that baits containing that second-generation anticoagulant at 0.005% active ingredient (a.i.) are likely to be effective against Norway rats and house mice of United States origin that are resistant to warfarin and other first generation anticoagulants. Data submitted to the Agency suggest that bromadiolone baits at 0.005% a.i. are likely to control warfarin-resistant Norway rats of United States origin and that 0.0025% difethialone baits that are registered for controlling house mice

are likely to control warfarin-resistant house mice of United States origin. The expression “warfarin-resistant” is used here because the test subjects ultimately killed by the second-generation anticoagulants mentioned initially were screened for resistance to warfarin.

Some success in controlling United States rodent populations resistant to warfarin has been reported in field trials involving second-generation anticoagulants and bromethalin (Ashton, *et al*, 1983). Anticoagulant-resistant rodents likely would be susceptible to any rodenticide that is not an anticoagulant. Alternating anticoagulant use with use of alternative control methods, including a toxicant with a different mode of action, would be expected to lessen the selective pressure favoring anticoagulant-resistant individuals. Frantz and Madigan (1998) suggested that a non-chemical strategy should be considered in areas where resistant rodents exist and that habitat modifications, especially limiting alternate food sources, might enhance the effectiveness of cyclical applications of first-generation anticoagulants in such areas by increasing the amount of bait consumed. Frantz and Madigan (1998) and Baroch (2004) present data which suggest that, if used cyclically and in conjunction with sanitation, Warfarin itself may eventually control resistant populations.

**Conditioned Food Aversion (Bait Shyness)** - Animals that become ill after ingesting a new food with a novel taste often avoid that food taste subsequently. When the new food is a toxic rodenticide bait, rodents obtaining sublethal oral doses may avoid both the bait base material and the toxicant in the bait base in the future, perhaps for the rest of their lives. Such acquired bait shyness can greatly diminish the effectiveness of rodent control programs and is most pronounced with quick-acting poisons that have salient flavors. Most such compounds no longer are registered in the United States. Among the rodenticides considered here, bait shyness is often a significant problem with zinc phosphide. With small target rodents such as voles (*Microtus* spp.), bait shyness to zinc phosphide can be mitigated appreciably by formulating baits to an active ingredient strength such that a single ingested bait particle is fatal to the rodent (Eadie, 1950). For larger species, prebaiting target mammals by exposing them to untreated bait grain several days before the toxic bait is applied may result in good control (Tietjen, 1976).

Due, at least in part, to the delayed toxic action of anticoagulants and cholecalciferol, bait shyness does not develop with these rodenticides. Bromethalin produces anorexia as an early symptom of exposure, but animals that recover do not seem to avoid the same bait subsequently.

**Weather** - Changes in ambient weather conditions can alter the behavior of rodents and may effect the amount of time that some species spend foraging above ground. Wet weather can also affect exposed baits directly by changing their integrity (especially with some types of pelleted baits) and by altering their toxic properties. Wet zinc phosphide baits “prematurely” liberate phosphine gas, reducing the toxicity of each particle and affecting palatability.

**Structural Use Considerations** - All rodenticide baits under discussion currently are registered for use to control commensal rodents in and around buildings. Consequently, each of these compounds

potentially is a viable alternative to each of the others in that use situation, except perhaps at sites where a high incidence of resistance to first-generation anticoagulants has been detected. Absent resistance problems, selection among active ingredients by professional applicators is likely to be based upon past experience, safety, cost, and possibly product promotions from manufacturers and/or distributors. In structural situations, most incidents for primary exposure are reported for young children (under 6 years of age) and for nontarget species, such as dogs, cats, and livestock, although primary exposures of wildlife may also occur. By following label directions regarding bait stations and proper use of tamper-resistant bait stations, professional users can minimize incidents of primary exposures to nontarget species. As secondary exposures are not affected by the use of bait stations and may occur even if all bait placements are located indoors, professional users may consider the likelihood of secondary exposures of avian and mammalian predators and scavengers when selecting among rodenticides.

A large proportion of rodenticide use by non-professional applicators (i.e., homeowners) occurs in structural situations, chiefly in and around their homes. Selection among rodenticide products by consumers depends to a large extent on local availability. The labels on rodenticide baits that consumers purchase are similar to the labels on baits used by professional applicators, but tamper-resistant bait stations are seldom offered at locations where consumers buy rodenticides. Some ready-to-use bait station products (bait stations loaded with bait) are offered for sale at retail outlets, but none of those stations has been demonstrated to be tamper-resistant. Ready-to-use bait stations that are made of cardboard are not tamper-resistant.

**Field Uses** - Most field uses of rodenticide bait are, or are soon to be, limited to labels of products that are classified as "Restricted Use Pesticides". Consequently, most applications to control field rodents or commensal rodents in field situations are to be made by certified applicators or by those under their direct supervision. Such applicators are expected to have access to training materials and equipment appropriate for the types of applications that they intend to make. Broadly, such types of applications include: above-ground treatments made manually or by aerial or ground broadcast equipment or by trail-builders; and below-ground applications made by hand, by hand-operated probes, or by machine-drawn burrow builders. Manual, above-ground treatments include "spot" placements of small amounts of bait in discrete locations, scattering bait over small areas near active burrows or runways, and placing larger amounts of bait in bait stations.

All above-ground placements are available to nontarget species to some extent. For spot, scattering, and trail-builder applications, some mitigation of primary nontarget exposure is realized by confining treatments to places where target species are likely to be the first animals to encounter baits. Bait stations used in the control of field rodents often are designed to exclude nontarget animals that are larger than the target species. In some cases, special designs have been developed which exploit behavioral traits and limitations of species of small nontarget mammals that are to be protected and allow control of larger rodents with little risk to those smaller species (e.g., Erickson , *et al*, 1990; Whisson, 1998)

With broadcast applications, spatial dilution of bait and its settling into vegetation may make it somewhat more likely to be encountered by the olfactory-guided foraging strategies of rodents than by visually guided granivorous birds.

For all above ground uses, proper timing of application in terms of season and time of day increases the likelihood of uptake by target species and thereby limits the amount of bait remaining for nontarget species.

Currently, for zinc phosphide, and following product reregistration for chlorophacinone and diphacinone, the field rodenticide uses available to nonprofessional applicators will be limited to manual, below-ground applications to control moles and/or pocket gophers. If label directions for such products are followed, primary nontarget exposures to such baits would be limited to species that entered the burrow systems of the target species. The likelihood that consumers would follow such directions may be greater for the mole and pocket gopher baits than for baits used to control commensal rodents because it is not necessary to obtain additional equipment to treat mole and gopher burrows and because there is little hope for successful control of moles and pocket gophers unless bait is applied directly to their burrow systems.

**Quarantine Uses** - When there is a need to ensure that no target pests survive, the selection of a rodenticide or combination of rodenticides is likely to favor the most highly toxic compounds. Whether there is a need to control a vector or an invasive rodent, a short-term increase in local risk factors might be offset by the long-term benefits of arresting the disease or preventing the invasive species from establishing itself.

**Conservation Uses** - If an invasive rodent species has become established, rodenticide baits may be incorporated in management or eradication programs intended to benefit native species. Such programs may be feasible on islands of small to moderate size to protect colonies of ground- and hole-nesting birds. If eradication is judged to be feasible, one season's use of a highly toxic rodenticide could eliminate the invasive rodent population to the subsequent benefit of the native species. Applying baits after migratory species have left the area may simultaneously promote acceptance of bait by food-stressed rodents while minimizing adverse impacts on native species. Some adverse impacts of island baiting programs on native species have been documented (e.g., Howald, *et al*, 1999). Recognizing such potential impacts and planning to mitigate are essential steps in designing strategies for managing or eradicating rodents for conservation purposes (e.g., National Park Service, 2000).

Use of rodenticide baits for conservation is mainly limited to government agencies.

## VIII. Alternative Methods of Control

### A. Chemical Alternatives

**Fumigants** (gas cartridges, aluminum phosphide, magnesium phosphide, Vargon, acrolein [SLN products only]) - Some types of rodents can be controlled by application of fumigant products to their burrow systems. Such applications potentially will kill target species and any other animals in the burrows. However, burrow treatments with some fumigants have been less than fully effective for organisms such as pocket gophers and moles, which have complicated burrow systems and may also have behavioral defenses against certain types of fumigants. Apparently for these reasons, gas cartridges do not appear to be effective against pocket gophers (e.g., Matschke, *et al* 1995).

The usefulness of burrow fumigants is further limited by prohibitions on treatments of burrows that are close to buildings and the need for soil moisture to be adequate to retain toxic gases. Especially under dry conditions, using gas cartridges may present a fire hazard. Aluminum phosphide, magnesium phosphide, and acrolein are Restricted Use Pesticides for which applicator certification and specialized training are required. Special equipment is needed to apply acrolein and Vargon to burrows. Burrow fumigations are time-consuming and labor-intensive, with repeat treatments often being needed. Consequently, burrow fumigations may be too costly to conduct except when controlling small infestations, "mopping up" survivors from a toxic baiting program, or protecting extremely valuable resources.

Structural fumigation aimed at controlling insect pests may also control exposed rodents. Benefits realized in this regard usually are in addition to the invertebrate control which was the primary treatment objective.

**Repellents** (e.g., Dr. T's., Ro•Pel, Naphthalene) - There are few products registered in the United States as rodent repellents. Such products generally are claimed to be area repellents or feeding deterrents. Evidence supporting either type of claim is limited. Denatonium compounds have been claimed to act as feeding deterrents to certain types of rodents as well as other mammals. At lower concentrations (e.g., 0.001%), however, denatonium benzoate is used in rodenticide baits and is claimed not to deter feeding by commensal rodents (Kaukeinen and Buckle, 1992).

Area repellents containing materials such as naphthalene or paradichlorobenzene are irritating to humans and are not appropriate for use in occupied buildings unless the products themselves are in sealed containers, in which case they would be useless as rodent repellents.

## B. Non-Chemical Alternatives

**Exclusion and Rodentproofing** - Rodent-resistant and rodentproof modifications to existing or new constructions (collectively called “stoppage”) can greatly reduce the likelihood that rodents will enter buildings and are an important element of successful structural rodent control operations.

However, not all building designs allow for complete rodent proofing, considering that young commensal rats can enter holes ½" (13 mm) in diameter and house mice can enter even narrower openings. 1988; (Baker, *et al*, 1994). Even with conventional rodent proofing, rats can occasionally enter buildings by coming up through the plumbing that drains toilets. Business and farm operations which require doors to be kept open for extended periods of time, especially at night, cannot be completely rodentproofed. Exclusion is not useful as a first approach to existing infestations as the immediate problem is to control the rodents.

**Habitat Modification** - Properties can be made less attractive to commensal rodents by rather simple procedures such as improving sanitation, thereby reducing sources of harborage, food, and water (Timm, 1994a; 1994b). These approaches are helpful in deterring colonization by rodents and, as directed on existing labels for commensal rodenticides, are useful follow-up practices after a successful rodent control program

**Trapping** - Trapping is useful for verifying the species responsible for a rodent infestation and may be sufficient to control small infestations, especially at the time of colonization (e.g., late summer invasions of homes by house mice). For larger infestations, rodent removal by trapping alone can be an extremely labor-intensive and costly (Boddicker, 1983). Where toxic baits cannot be used for reasons of safety, trapping may be the only removal method that can be implemented. Lethal and non-lethal traps of various types may be used to control various types of rodents and moles. Traps may be especially useful for controlling small infestations of house mice, *Peromyscus* spp. mice, pocket gophers, woodchucks, and Norway rats. Appropriate trap placement is important to success, and trap prebaiting may enhance the rate of capture of neophobic species (e.g., Timm, 1994a; 1994b).

Lethal traps also pose a risk to small nontarget animals, and can cause injury to applicators and other on-site personnel. Placing lethal traps within locked rooms or in the rodent areas of properly applied tamper-resistant bait stations or other suitable objects can mitigate the risks posed to some nontarget species.

Live-trapping requires users to deal with captured animals. This circumstance often allows for the release of captured nontarget animals but presents to the trapper the problem of deciding the fate of target animals and may increase the opportunity for transfer of certain disease to humans (e.g., CDC, 2004). With repeating-type mouse traps, multiple house mice may be caught in a single trap (Timm, 1994a). With traps, there is some danger of targeted animals becoming trap-shy through their own experiences (i.e., near misses).

Electrical or electronic rodent traps have been marketed at times. These traps generally have been able to trap rodents, with some traps also killing them. Due to high unit costs, it is doubtful that such traps have had much usage.

**Shooting** - Some control of diurnal rodents such as prairie dogs and ground squirrels can be effected through shooting. The method has been described as selective, but not very efficient for controlling colonial species (e.g., Hyingstrom and Virchow, 1994; Marsh, 1994). For woodchucks, shooting can be used to manage local populations (Bollengier, 1994). However, shooting can only be employed where and when State laws and local ordinances permit the activity. Prohibitions on hunting near buildings or in parks and the “game” status of the pest are some of the factors affected by laws and ordinances.

**Pest Control Devices** - Under FIFRA, pest control devices are regulated to some extent, but there is no requirement that the products themselves be registered before they are marketed. Some pest control devices marketed for rodent control are discussed below.

**Glue boards** - Glue boards are sticky traps which have been marketed in ready-to-use condition or with the glue offered separately, enabling the user to apply it to a suitable surface for capturing rodents. The issues with glue boards are much like those discussed above for traps, except for logistical and humaneness problems since the glue restrains but does not kill the rodent, often leading to prolonged struggling in awkward positions (Frantz and Padula, 1983). Humans may be injured when picking up glue boards holding rodents that only appear to be dead. Neophobic responses to glue boards have been reported for house mice (Corrigan, 1998).

**Burrow collapsers** - Products that collapse rodent burrow systems are relatively new. One type of such a device ignites a mixture of propane and oxygen in underground burrow systems, causing an explosion which reportedly kills organisms living in the burrow. Compression and the collapsing of the tunnels are believed to effect rodent control, with the leveled burrow system subsequently being less inviting to immigrant rodents than the intact system would have been. The utility of this type of product would be limited to burrowing species.

Issues of safety to humans, property, environment are largely unstudied for this type of product. Appropriate personal protective equipment (PPE) should be worn when using a burrow collapsing device. Nontarget animals living in treated burrow systems also would be expected to be killed. The abrupt shifting of soil along the course of the treated burrow system could cause damage to nearby buildings, roads, fencing, utility poles, or other structures.

**Electromagnetic Devices** - Since the mid 1970's, various devices have been claimed to be able to control various types of pests, including rodents, through creation of electromagnetic fields. EPA collected and evaluated many such units that were available in the 1970's and concluded that the principle of controlling pests via low levels of electromagnetism is flawed (EPA, 1979). Consequently,

electromagnetic devices are considered to be unlikely to exert rodent control effects on their own; and claims that they enhance the effectiveness of conventional rodent control techniques such as traps and toxic baits are regarded as highly unlikely.

**Ultrasonic devices** - Although many types of rodents can perceive ultrasonic stimuli and communicate via ultrasonic vocalizations, tests of ultrasound generators as rodent control agents have suggested little promise for them. Even though rodents may appear to perceive and may initially react to ultrasonic noise, they seem to adapt to it. Ultrasound rapidly attenuates over distance, is extremely directional, and is “shadowed” by objects. Consequently, commensal rodents appear to be able to coexist with ultrasonic signals by staying largely behind and under cover, which is typical behavior for these animals. EPA studied units available in the 1970's and early 1980s and concluded that they were not likely to be of significant value in rodent control.

### **C. The Integrated Pest Management (IPM) Approach**

Management of rodent pests generally is best achieved through a combination of viable methods (i.e., IPM) , with the approaches selected being adapted to conditions prevailing at the site when the program is initiated. For example, Timm (1994) presents a flow chart outlining the factors and control options that an applicator might consider in dealing with an infestation of house mice. If there are no rodents present, rodentproofing and habitat modification should be implemented to increase the likelihood that rodents will not take up residence on the premises. Planning to incorporate rodent stoppage into building designs, seeing to it that stoppage is effected during construction, and diligently maintaining stoppage and sanitation on the premises may prevent rodent infestations from occurring at a site (Frantz and Davis, 1991).

Frantz (2004) considers the following criteria for selecting a rodenticide for use in an IPM programs for rodents, especially one involving public health: a) the rodenticide should be the least toxic product that will be effective on the target species, and b) the rodenticide must have a highly efficacious and readily available antidote that typically can be administered in time to save an accidentally intoxicated animal or human.

For centuries, however, there have been rodent infestations in many types of habitat on all continents and on most islands around the world. Consequently, there will continue to be many occasions in which lethal control methods for rodents will be sought; and rodenticide baits will continue to be needed for the foreseeable future.

## **IX. Matrix of Use Sites**

Table 1. summarizes the use patterns of the nine rodenticides discussed here. The names of the rodenticides are listed on the first column. The second column shows a common classification scheme for these chemicals based on mode of action and developmental history. The next two columns

*September 2, 2004*

indicate that all nine rodenticides are registered for commensal rodent control, whereas only four of these, warfarin, chlorophacinone, diphacinone, and zinc phosphide are registered for non-commensal rodent control. Of these, zinc phosphide has by far the most uses in terms of number of target species and use sites. The fifth column shows the ranking of the four rodenticides that pose the greatest overall risk to birds and nontarget mammals, based on EPA's 2004 ecological risk assessment for the nine rodenticides discussed in this document. The sixth column shows how the nine rodenticides are rated in terms of IPM-compatibility according to Frantz (2004), who uses relative toxicity and antidote availability for poisoned nontarget animals as criteria for establishing the IPM-compatibility of rodenticides a rodent management expert. This column is included to stimulate public comments and further discussion. The next to the last column includes some of the limited usage data publicly available at present. The last column attempts to categorize common use patterns in relation to the areas where these rodenticides are most commonly used.

September 2, 2004

**Table 1. Use Matrix for Currently Registered Rodenticide Baits**

Rodenticide Baits	Active Ingredient Classification	Use Against Common Pest Mammals		Potential Overall Risk to Birds and Nontarget Mammals <sup>4</sup>	“IPM-Compatible” <sup>5</sup>	Usage <sup>6</sup>	Land Use Category (Use is Mainly in these Areas)
		Commensal Rats and Mice	Mainly Native Rodents <sup>2</sup>				
Warfarin <sup>1</sup>	1 <sup>st</sup> Generation Anticoagulant	X	X		Yes	No Estimate Available	Urban / Suburban
Chlorophacinone	1 <sup>st</sup> Generation Anticoagulant	X	X (Sect. 24c) (RUP <sup>3</sup> )		Yes	No Estimate Available	Urban / Suburban / Rural
Diphacinone	1 <sup>st</sup> Generation Anticoagulant	X	X (Sect. 24c) (RUP <sup>3</sup> )	High (4 <sup>th</sup> )	No	No Estimate Available	Urban / Suburban / Rural
Brodifacoum	2 <sup>nd</sup> Generation Anticoagulant	X		High (1 <sup>st</sup> )	No	Most used homeowner rodenticide. Products have an estimated 30% of PCO market	Urban / Suburban
Bromadiolone	2 <sup>nd</sup> Generation Anticoagulant	X			Yes <sup>7</sup>	Products have an estimated 50% of PCO market	Urban / Suburban
Difethialone	2 <sup>nd</sup> Generation Anticoagulant	X		High (3 <sup>rd</sup> )	No	No estimate Available	Urban / Suburban
Zinc phosphide	Non-Anticoagulant	X	X (Sect. 3) (RUP <sup>3</sup> )	High (2 <sup>nd</sup> )	Yes <sup>7</sup>	Most used rodenticide in agricultural fields	Rural / Agriculture
Cholecalciferol	Non-Anticoagulant	X			Probably not	No estimate Available	Urban / Suburban
Bromethalin	Non-Anticoagulant	X			No	No estimate Available	Urban / Suburban

<sup>1</sup> Resistance to this a.i. has been reported in some United States commensal rat populations.

<sup>2</sup> Also jackrabbits, moles, mongoose, and nutria (an introduced rodent)

<sup>3</sup> Restricted Use, except for underground baiting of pocket gophers and moles

<sup>4</sup> Based on Erickson and Urban (2004)

<sup>5</sup> Based on Frantz (2004), who uses relative toxicity and antidote availability for poisoned nontarget animals as criteria for establishing the IPM-compatibility of rodenticides.

<sup>6</sup> There is a limited amount of data available on the usage of rodenticides in the three primary rodenticide markets (homeowner, PCO (residential and commercial), and agriculture). Not Available means that there is no estimate available for the usage of the rodenticide. Not Available does not mean that the rodenticide is not used. Annual value of homeowner market for rodent control estimated at more than \$90 million annually. The estimated annual value of the market for rodent control products used by pest control operators (PCOs) is more than \$15 million.

<sup>7</sup> Only when there is a critical need to rapidly knock down a rodent population as, for instance, when facing a rodent-borne disease outbreak.

## **X. Topics for Specific Public Comment**

As part of the public comments process, the Agency is interested in obtaining additional information related to topics where data are limited or where more current information on topics addressed is available. Examples of reliable information needed include the following:

- Prevalence of resistance to baits containing warfarin or any other rodenticide in any target rodent, use site, or geographic area, especially in the United States
- Examples of how resistance to rodenticide baits have been successfully managed.
- Examples of successful commensal rodent control programs in urban areas.
- Examples of successful integrated pest management (IPM) programs targeting any type of rodent pests.
- Amounts of annual use for all nine rodenticide baits discussed in this document.
- Comparative efficacy data for rodenticide baits.
- Wholesale and retail price of rodenticide baits.
- Relative importance of rodenticide baits relative to nonchemical control methods.
- Common zinc phosphide uses, including main target pests and use sites.
- Detailed estimates of the types of damage caused by rodents in the United States and the economic loss resulting from such damage.

## **XI. References**

Albert, S.W. and C.R. Record. 1979. Efficacy and cost of four rodenticides for Columbian ground squirrel control. Unpublished report: Montana Department of Livestock, Helena, MT, 22 pp.

Andrews, J.M. and Link, V.B. 1947. The murine typhus fever problem in the United States. *Pests and Their Control*, January issue.

Ashton, D.A., W.B. Jackson, and J.H. McCumber. 1983. An evaluation of methods used in

September 2, 2004

- comparative field testing of commensal rodenticides. Pages 138 - 154 In: Kaukeinen, D.E. (Ed.) Vertebrate Pest Control and Management Materials: Fourth Symposium, ASTM Special Technical Publication 817, American Society for Testing and Materials, Philadelphia, PA.
- Baker, R.O., R.M. Timm, and G.R. Bodman. 1994. Rodentproof construction and exclusion methods. Pages B-137 - B-150 In: Hyngstrom, S.E., Timm, R.M., and Larson, G.E. (Eds.) Prevention and Control of Wildlife Damage, University of Nebraska Cooperative Extension, Institute of Agriculture and Natural Resources, University of Nebraska, Lincoln, NE; and U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Animal Damage Control, Riverdale, MD.
- Baroch, J. (2004) Warfarin: a forgotten weapon? *Pest Control Technology*, August issue, 44-52.
- Beall, J.C. 1946. Permanent program for typhus fever control on Memphis, Tennessee. *American Journal of Public Health*, 36(2):161-168.
- Blindauer, K. 1999. Rodenticide benefits: A public health perspective, Centers for Disease Control and Prevention (CDC), National Center for Environmental Health, Presentation at the June 9, 1999 Rodenticide Stakeholder Workgroup,
- Boddicker, M.L. 1983. Prairie dogs. In: Timm, R.M. (Ed) Prevention and Control of Wildlife Damage, Great Plains Agricultural Council Wildlife Resources Committee and Nebraska Cooperative Extension Service Institute of Agriculture and Natural Resources, University of Nebraska, Lincoln, NE, B-75 to B-84.
- Bollengier, R.M. Jr. 1994. Woodchucks. Pages B-183 - B-187 In: Hyngstrom, S.E., Timm, R.M., and Larson, G.E. (Eds.) Prevention and Control of Wildlife Damage, University of Nebraska Cooperative Extension, Institute of Agriculture and Natural Resources, University of Nebraska, Lincoln, NE; and U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Animal Damage Control, Riverdale, MD.
- Boren, J. and R. Valdez. 2003. Rodent control and protection from hantavirus. New Mexico State University, Cooperative Extension Service, Guide L-209.  
<http://www.cahe.nmsu.edu/pubs/1/1-209.pdf>
- Brooks, J.E. and L.A. Fiedler. 1999. Compendium of Post-Harvest Operations. Chapter III : Vertebrate Pests: Damage on Stored Foods, Agro-Industries and Post-Harvest Management Service, Food and Agriculture Organization (AGSI-FAO), Rome.  
<http://www.fao.org/inpho/compend/text/ch03.htm>

- Brown, S.L. 1994. Insect management in grain stored on the farm: Importance of stored grain pests. The University of Georgia, College of Agricultural & Environmental Sciences, Cooperative Extension Service, Bulletin 1101/1994.  
<http://www.ces.uga.edu/pubcd/b1101-w.html>
- CDC. 1998. Rat-bite fever -- New Mexico, 1996. Morbidity and Mortality Weekly Report, Center for Disease Control and Prevention, 47(05):89-91.  
<http://www.cdc.gov/epo/mmwr/preview/mmwrhtml/00051368.htm>
- CDC. 1999. All about hantaviruses. National Center for Infectious Diseases, Special Pathogens Branch, Center for Disease Control and Prevention.  
<http://www.cdc.gov/ncidod/diseases/hanta/hps/noframes/teach.htm>
- CDC. 2002. Murine typhus -- Hawaii, 2002. Morbidity and Mortality Weekly Report, Center for Disease Control and Prevention, 52(50):1224-1226.  
<http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5250a2.htm>
- CDC. 2003. Fact Sheet: Trichinellosis, Division of Parasitic Diseases, Center for Disease Control and Prevention.  
[http://www.cdc.gov/ncidod/dpd/parasites/trichinosis/factsht\\_trichinosis.htm](http://www.cdc.gov/ncidod/dpd/parasites/trichinosis/factsht_trichinosis.htm)
- CDC. 2003. Information on plague. Division of Vector-Borne Infectious Diseases, Center for Disease Control and Prevention. <http://www.cdc.gov/ncidod/dvbid/plague/info.htm>
- CDC. 2003. Leptospirosis. Division of Bacterial and Mycotic Diseases, Center for Disease Control and Prevention. [http://www.cdc.gov/ncidod/dbmd/diseaseinfo/leptospirosis\\_g.htm](http://www.cdc.gov/ncidod/dbmd/diseaseinfo/leptospirosis_g.htm)
- CDC. 2004. Hantavirus pulmonary syndrome: Trap up! Center for Disease Control and Prevention, Atlanta, GA, April 26, 2004, 1 p  
[www.cdc.gov/ncidod/diseases/hanta/hps\\_stc/stc\\_trap.htm](http://www.cdc.gov/ncidod/diseases/hanta/hps_stc/stc_trap.htm)
- Cornell University. 1997. New forms of old disease, leptospirosis, threaten dogs in U.S., Cornell veterinarians warn humans also are susceptible to flu-like infection from urine of rodents, cattle, other animals. Science News, October 7, 1997.  
<http://www.news.cornell.edu/releases/Oct97/lepto.hrs.html>
- Cornell University. 2001. Zinc phosphide (Phosvin, KP) Chemical Profile 1/85.  
<http://pmep.cce.cornell.edu/profiles/rodent/zinc-phosphide/rod-prof-zinc-phosphide.html>

- Corrigan, R.M. 1998. The efficacy of glue traps against wild populations of house mice, *Mus domesticus*, Ruttly. Pages 268-275 In: Baker, R.O. and Crabb, A.C. (Eds.), *Proceedings: Eighteenth Vertebrate Pest Conference*, 18, University of California at Davis.
- Dent, J.E., Morgan, H.B., and Hill, E.L. 1949. Effects of DDT dusting on domestic rats under colony and field conditions. *Public Health Reports*, 64(21):666-671.
- Department of Environmental Health, County of San Diego. Undated. Vector Surveillance and Control: Diseases Transmitted by Domestic Rodents.  
<http://www.co.san-diego.ca.us/deh/chd/vector/vratdiseas.html>
- Donlan, J.C., G.R. Howald, B.R. Tershy, and D.A. Croll. 2003. Evaluating alternative rodenticides for island conservation: roof rat eradication from the San Jorge Islands, Mexico. *Biological Conservation* 114(2003):29-34
- Eadie, W.R. 1950. New techniques in control of orchard mice. *Cornell University Agricultural Experiment Station Bulletin No. 856*. 15 pp.
- Emlen, J.T. and A. W. Stokes. 1947. Effectiveness of various rodenticides on populations of brown rats in Baltimore, Maryland. *American Journal of Hygiene*, 43:254-257.
- Erickson, W.A., R.E. Marsh, and W.L. Halvorson. 1990. A roof rat bait station that excludes deer mice. *Wildlife Society Bulletin*, 18:319-325.
- Erickson, W. and D. Urban. 2004. Potential risks of nine rodenticides to birds and nontarget mammals: a comparative approach, Final Assessment. U.S. Environmental Protection Agency.
- Ebeling, W. 1975. Urban Entomology. University of California, Division of Agricultural Sciences. 695 pp.
- Emlen, J.T. and A.W. Stokes. 1947. Effectiveness of various rodenticides on populations of brown rats in Baltimore, Maryland. *American Journal of Hygiene*, 43:254-257.
- Fisher, D.D. and S.E. Hygnstrom. 1997. Controlling vole damage, NebGuide, University of Nebraska, Lincoln. <http://ianrpubs.unl.edu/wildlife/g887.htm>
- Frantz, S.C. 1988. Architecture and commensal vertebrate pest management. Pages 228-295 In R.B. Kundsinn (Ed.), *Architectural design and indoor microbial pollution*. Oxford University Press, New York, NY

September 2, 2004

- Frantz, S.C. 2004. Communication to Jerry Hershovitz. Center for Disease Control and Prevention, National Center for Environmental Health, Emergency and Environmental Health Services.
- Frantz, S.C. and Davis, D.E. (1991) Bionomics and integrated pest management of commensal rodents. In: Gorham, J.R. (ed.) Ecology and Management of Food-Industry Pests, FDA Technical Bulletin 4, Association of Official Analytical Chemists, Arlington, VA, 243-313.
- Frantz, S.C. and C.M. Padula. 1980. Recent developments in anticoagulant resistance studies: surveillance and application in the U.S. Pages 80-88 In: Clark, J.P. and Marsh, R.E. (Eds.) Proceedings: Ninth Vertebrate Pest Conference, 9, University of California at Davis, CA.
- Frantz, S.C. and C.M. Padula. 1983. A laboratory test method for evaluating the efficacy of glueboards for trapping house mice. Pages 209-225 In D.E. Kaukeinen (Ed.) Vertebrate Pest Control and Management Materials: Fourth Symposium, ASTM STP 817, American Society for Testing and Materials, Philadelphia, PA.
- Frantz, S.C. and C.P. Madigan. 1998. Warfarin resistance revisited. Pages 268-275 In: Baker, R.O. and Crabb, A.C. (Eds.), Proceedings: Eighteenth Vertebrate Pest Conference, 18, University of California at Davis, CA.
- Frieden, T.R. 2003. Testimony, New York City Department of Health and Mental Hygiene, Rodent Control in New York City, Testimony Before New York City Council, Committees on Health and on Governmental Operations, October 28, 2003, City Hall, New York, New York. <http://www.nyc.gov/html/doh/html/public/testi/testi20031028.html>
- Harrington, E.P., P.E. Kaufman, D.B. Weingart, J.K. Walfron, and D.A. Rutz. 1998. Pest and pesticide use assessment for poultry production systems in New York State for 1998. Pesticide Management Education Program, Cornell University, Ithaca, NY. <http://pmep.cce.cornell.edu/piap/poultry98/index.html>
- Howald, G.R., Mineau, P., Elliott, J.E., and Cheng, K.M. (1999) Brodifacoum poisoning of avian scavengers during rat control on a seabird colony. *Ecotoxicology* 8:431-447.
- Hayes, W.J. Jr. and T.B. Gaines. 1950. Control of rats with residual rodenticide Warfarin. *Public Health Reports*, 65(47):1537-1555.
- Hill, E.L. and Morlan, H.B. 1948. Evaluation of county-wide DDT dusting operations in murine typhus control. *Public Health Reports*, 68(51):1635-1653.

- Hopf, H.S., G.E. Morley, and J.E. Humphries. 1976. Rodent damage to growing crops and to farm and village storage in tropical and subtropical regions. pp. 115. *Cent. Overseas Pest Res.*, London, UK.
- Hyngstrom, S.E. 1995. Vertebrate pest damage in grain storage facilities. Pp. 227-238 *In: Stored Product Management*, Oklahoma Cooperative Extension Service, U.S. Department of Agriculture, Federal Grain Inspection Service, Animal and Plant Health Inspection Service, Circular No. E-912.
- Hyngstrom, S.E. and D.R. Virchow. 1994. Prairie dogs. Pages B-151 - B-158 *In: Hyngstrom, S.E., Timm, R.M., and Larson, G.E. (Eds.) Prevention and Control of Wildlife Damage*, University of Nebraska Cooperative Extension, Institute of Agriculture and Natural Resources, University of Nebraska, Lincoln, NE; and U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Animal Damage Control, Riverdale, MD.
- Kaukeinen, D.E. and A.P. Buckle. 1992. Evaluation of aversive agents to increase the selectivity of rodenticides with emphasis on denatonium benzoate (Bitrex) bittering agent. Pages 192-199 *In: Borrecco, J.E. and Marsh, R.E. (Eds.) Proceedings: 15th Vertebrate Pest Conference*, 15, University of California at Davis, CA.
- Larson, E. and Morgenthaler, E. 1994. Killer disease, borne by rodents, is found in wider areas of U.S. *Wall Street Journal*, 223(10):A1-A2.
- Link, K.P. 1959. The discovery of dicumarol and its sequels. *Circulation*, 19(1):97-107.
- Marsh, R.E. 1994. Belding's, California, and rock ground squirrels. Pages B-151 - B-158 *In: Hyngstrom, S.E., Timm, R.M., and Larson, G.E. (Eds.) Prevention and Control of Wildlife Damage*, University of Nebraska Cooperative Extension, Institute of Agriculture and Natural Resources, University of Nebraska, Lincoln, NE; and U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Animal Damage Control, Riverdale, MD.
- Marsh, R.E., Howard, W.E., and Jackson, W.B. (1980) Bromadiolone: a new toxicant for rodent control for rodent control. *Pest Control*, 48, 22-26.
- Matschke, G.H., Ramey, C.A., McCann, G.R., and Engemann, R.M. 1995. Evaluation of a 2-active ingredient gas cartridge for controlling Northern pocket gophers. *International Biodeterioration & Biodegradation*, 36(1/2):151-160.

September 2, 2004

- Mills, J.N., A. Corneli, J.C. Young, L.E. Garrison, A.S. Khan, and T.G. Ksiazek. 2002. Hantavirus pulmonary syndrome - United States: updated recommendations for risk reduction, CDC, *Morbidity and Mortality Weekly Report*, 51: RR-9, July 26, 2002. <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5109a1.htm>
- Mohr, C.O. and Smith, W.W. 1957. Eradication of murine typhus fever in a rural area. *Bulletin of the World Health Organization*, 16:255-266.
- Montana Department of Agriculture. undated. Voles in Montana: their biology, damage and control. <http://agr.state.mt.us/programs/asd/vertpest.shtml>
- Montana Department of Agriculture. undated. Pocket Gopher Control Techniques. <http://agr.state.mt.us/programs/asd/vertpest.shtml>
- Morlan, H.B. and Hines, V.D. 1951. Evaluation of county-wide DDT dusting operations in murine typhus control, 1950. *Public Health Reports*, 66(33):1052-1057.
- National Park Service (2000) *Anacapa Island Restoration Project*, Final Environmental Impact Assessment, U.S. Department of the Interior, Channel Islands National Park, Ventura, CA, 140 pp.
- NPCA 1994. Hantavirus. Technical Release ESPC 071939 3/94, National Pest Control association, Dunn Loring, VA, 4 pp.
- Pinto, L. 1992. Urban integrated pest management, a guide for commercial applicators. Module III, Vertebrates. Pinto & Associates, Inc. Prepared for the United States Environmental Protection Agency.
- Pratt, H.D., B.F. Bjornson, and K.S. Litting. 1976. Control of domestic rats & mice. U.S. Department of Health and Human Services, Public Health Services, Center for Disease Control, Atlanta, Georgia, HHS Publication No. (CDC) 81-8141.
- Proctor, D.L. 1994. Grain storage techniques: evolution and trends in developing countries. FAO, Agricultural Services Bulletin No. 109. <http://www.fao.org/docrep/T1838E/T1838E00.htm#Contents>
- Salmon, T.P. and W.P. Gorenzel. 2002. Voles (Meadow Mice). University of California Pest Management Guidelines. <http://www.ipm.ucdavis.edu/PMG/PESTNOTES/pn7439.html>

September 2, 2004

- Schneider, B.A. and R.K. Hitch. 1982. *Pesticide Assessment Guidelines*, Subdivision G, Product Performance. Office of Pesticides and Toxic Substances, U.S. Environmental Protection Agency, 422 pp.
- Scott, H.G. (1982) A neglected technique in integrated pest control. *Pest Control*, July, 41-45.
- Schwartz, P. 2003. Bubonic plague, U.S. Department of the Interior, National Park Service, Public Health program. [http://www.nps.gov/public\\_health/inter/info/factsheets/fs\\_plague.htm](http://www.nps.gov/public_health/inter/info/factsheets/fs_plague.htm)
- State of Colorado Department of Agriculture. Undated. Rodent/predator control. <http://www.ag.state.co.us/animals/rodent/html>
- Tickes, B.R., L.K. Cheatham, and J.L. Stair. 1982. A comparison of selected rodenticides for the control of the common valley pocket gopher (*Thomomys bottae*). In: Marsh, R.E. (ed) *Proceedings: Tenth Vertebrate Pest Conference*, 10, University of California at Davis, CA, 201-204.
- Tietjen, H.P. 1976. Zinc phosphide -- its development as a control agent for black-tailed prairie dogs. Special Scientific Report -- Wildlife No. 195, U.S. Fish and Wildlife Service, U.S. Department of the Interior, Washington, DC, 18 pp.
- Tietjen, H.P. 1979. Zinc Phosphide -- its efficacy as a control agent for Gunnison (Zuni) prairie dogs (*Cynomys gunnisoni gunnisoni* and *C. g. zuniensis*). Completion report for research conducted under work units DF-924.07 and DF-924.07A (Amended), Denver Wildlife Research Center, Lakewood, CO, U.S. Department of the Interior, U.S. Fish and Wildlife Service, 16 pp.
- Tietjen, H.P. and G.H. Matschke. 1981. An evaluation of zinc phosphide for the management of black-tailed prairie dogs (*Cynomys ludovicianus*). 1981 Job Completion Report for Experimental Use Permit No. 6704-EUP-25, Shannon County, South Dakota. Denver Wildlife Research Center, U.S. Fish and Wildlife Service, U.S. Department of the Interior, 17 pp.
- Tietjen, H.P. 1976. Zinc Phosphide -- its development as a control agent for black-tailed prairie dogs. Special Scientific Report -- Wildlife No. 195, U.S. Fish and Wildlife Service, U.S. Department of the Interior, Washington, DC, 18 pp.
- Timm, R.M. 1994a. House mice. Pages B-31 - B-46 In: Hyngstrom, S.E., Timm, R.M., and Larson, G.E. (Eds.) *Prevention and Control of Wildlife Damage*, University of Nebraska Cooperative Extension, Institute of Agriculture and Natural Resources, University of Nebraska,

September 2, 2004

Lincoln, NE; and U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Animal Damage Control, Riverdale, MD.

Timm, R.M. 1994b. Norway rats. Pages B-105 - B-120 In: Hyngstrom, S.E., Timm, R.M., and Larson, G.E. (Eds.) Prevention and control of wildlife damage, University of Nebraska Cooperative Extension, Institute of Agriculture and Natural Resources, University of Nebraska, Lincoln, NE; and U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Animal Damage Control, Riverdale, MD.

Timm, R.M. and W.E. Howard. 1994. White-footed and deer mice, Prevention and control of wildlife damage. University of Nebraska - Lincoln, Cooperative Extension Division and USDA - APHIS - Animal Damage Control, B-47. <http://afsafety.af.mil/AFSC/Bash/USDA/rodent.html>

Townzen, K.R., Thompson, M.A., and Smith, C.R. 1996. Investigations and management of epizootic plague at Ice House Reservoir, Eldorado National Forest, California 1994 and 1995. Pages 68-74 In: Timm, R.M. and Crabb, A.C. (Eds.) Proceedings: 17th Vertebrate Pest Conference 17, University of California at Davis, CA.

USDA APHIS Veterinary Services. 1972. Facts About Trichinosis. <http://www.usaha.org/species/trichinosis.html#contents>

USDA NASS. 1999. Agricultural chemical usage: 1997 livestock and general farm survey, U.S. Department of Agriculture, National Agricultural Statistics Service.

U.S.EPA 1981. Warfarin and its sodium salt [3-(alpha-Acetylbenzyl)-4-hydroxycoumarin]. Pesticide Registration Standard, Office of Pesticides and Toxic Substances, U.S. Environmental Protection Agency, 133 pp.

U.S. EPA. 1998. Reregistration Eligibility Decision (RED): Zinc Phosphide, Office of Prevention, Pesticides and Toxic Substances.

U.S. EPA. 1979. EPA stops sale of several electromagnetic insect and rodent repellents. EPA "Environmental News" press release, April 2, 1979, 3 pp.

U.S. EPA. 2004. Proprietary data.

Whisson, D.A. 1998. Modified bait stations for California ground squirrel control in endangered kangaroo rat habitat. Pages 233-235 In: Baker, R.O. and Crabb, A.C. (Eds.), Proceedings: Eighteenth Vertebrate Pest Conference, 18, University of California at Davis, CA.

September 2, 2004

Wiley, J.S. 1946. Recent developments in murine typhus fever control. *American Journal of Public Health*, 36 (9):974-983.

## **XII. List of Contributors**

Angel Chiri, U.S. EPA, OPP, Biological and Economic Analysis Division

William Jacobs, U.S. EPA, OPP, Registration Division

Jonathan Becker, U.S. EPA, OPP, Biological and Economic Analysis Division

Tim Kiely, U.S. EPA, OPP, Biological and Economic Analysis Division

Kelly White, U.S. EPA, OPP, Special Review and Reregistration Division

Laura Parsons, U.S. EPA, OPP, Special Review and Reregistration Division

William Gross, U.S. EPA, OPP, Biological and Economic Analysis Division